



REVIEW ARTICLE

CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution

Murray R. Badger¹ and G. Dean Price

Molecular Plant Physiology Group, Research School of Biological Sciences, Australian National University,
PO Box 475, Canberra City, ACT 2601, Australia

Received 11 April 2002; Accepted 10 October 2002

Abstract

Cyanobacteria have evolved an extremely effective single-cell CO₂ concentrating mechanism (CCM). Recent molecular, biochemical and physiological studies have significantly extended current knowledge about the genes and protein components of this system and how they operate to elevate CO₂ around Rubisco during photosynthesis. The CCM components include at least four modes of active inorganic carbon uptake, including two bicarbonate transporters and two CO₂ uptake systems associated with the operation of specialized NDH-1 complexes. All these uptake systems serve to accumulate HCO₃⁻ in the cytosol of the cell, which is subsequently used by the Rubisco-containing carboxysome protein micro-compartment within the cell to elevate CO₂ around Rubisco. A specialized carbonic anhydrase is also generally present in this compartment. The recent availability of at least nine cyanobacterial genomes has made it possible to begin to undertake comparative genomics of the CCM in cyanobacteria. Analyses have revealed a number of surprising findings. Firstly, cyanobacteria have evolved two types of carboxysomes, correlated with the form of Rubisco present (Form 1A and 1B). Secondly, the two HCO₃⁻ and CO₂ transport systems are distributed variably, with some cyanobacteria (*Prochlorococcus marinus* species) appearing to lack CO₂ uptake systems entirely. Finally, there are multiple carbonic anhydrases in many cyanobacteria, but, surprisingly, several cyanobacterial genomes appear to lack any identifiable CA genes. A pathway for the evolution of CCM components is suggested.

Key words: Carboxysomes, CO₂ concentrating mechanisms, cyanobacteria, evolution, genes, photosynthesis, transporters.

Introduction

Cyanobacteria have existed as oxygenic photosynthetic bacteria on earth for at least 2.7 billion years (Buick, 1992). During that time they have endured a changing gaseous environment where CO₂ has declined and O₂ has risen. This has imposed evolutionary pressure on them to evolve strategies for efficiently acquiring inorganic carbon for photosynthesis. In response to this they have developed an effective photosynthetic CO₂ concentrating mechanism (CCM) for improving the carboxylation by their relatively inefficient Rubiscos (Badger and Price, 1992; Price *et al.*, 1998; Kaplan and Reinhold, 1999). This CCM is perhaps the most effective of any photosynthetic organism, concentrating CO₂ up to 1000-fold around the active site of Rubisco. In the past few years, there has been a rapid increase in the understanding of the mechanisms and genes involved in cyanobacterial CCMs. In addition, there has been a recent expansion in the availability of complete cyanobacterial genome sequences, thus increasing the potential to examine questions regarding both the evolution and diversity of components of the CCM across cyanobacterial species. This paper reviews current understanding of the mechanisms and genes underlying the operation of the cyanobacterial CCM, and takes the opportunity to employ comparative genomics to shed light on the evolution and diversity of the CCM among cyanobacterial species.

¹ To whom correspondence should be addressed. E-mail: murray.badger@anu.edu.au

Abbreviations: ABC, ATP binding cassette; CA, carbonic anhydrase; CCM, CO₂ concentrating mechanism; Ci, inorganic carbon; EZ, ethoxymethylamine; Fd, ferredoxin; NDH-1, NAD(P)H dehydrogenase; PQ, plastoquinone; Rubisco, ribulose 1–5, biphosphate carboxylase-oxygenase EC. 4.1.1.39.

The cyanobacterial CCM model

A schematic diagram of a cyanobacterial cell with components of the CCM (see recent reviews by Price *et al.*, 1998; Kaplan and Reinhold, 1999) is shown in Fig. 1. The basic concepts of this model are based on experiments with the model cyanobacterial species *Synechococcus* PCC7942, *Synechocystis* PCC6803 and *Synechococcus* PCC7002. Central to the functioning of the cyanobacterial CCM is the carboxysome, a protein micro-compartment within the cell that contains the Rubisco of the cell together with a carboxysomal carbonic anhydrase (CA). The CA functions to convert an accumulated cytosolic pool of HCO_3^- into CO_2 within the carboxysome. The generation of CO_2 coupled with a diffusive restriction to the efflux from the carboxysome, possibly imposed by the protein shell, leads to the localized elevation of CO_2 around the active site of Rubisco within the carboxysome. The substrate for the carboxysome, HCO_3^- , is accumulated in the cytosol by the operation of a number of active CO_2 and HCO_3^- transporters. These transporters are located on both the plasma membrane and the thylakoid membrane, and exist in both low affinity and high affinity transporter forms. Many cyanobacteria have the ability to improve their affinity for inorganic carbon (Ci) when grown at limiting




Fig. 1. A generalized model for the cyanobacterial CCM. Shown on the figure are the Rubisco-containing carboxysomes with the carboxysomal carbonic anhydrase (CA) and an associated diffusional resistance to CO_2 efflux. The accumulation of HCO_3^- in the cytosol is achieved through the action of a number of CO_2 and HCO_3^- uptake systems.

Ci levels, and this acclimation is due primarily to the changes that occur in the synthesis and properties of various high and low-affinity Ci transporters associated with the cells.

The phylogeny of cyanobacteria

Attempts to divide cyanobacteria into phylogenetic groupings have used both photosynthetic pigment composition and ribosomal RNA subunit sequences. The prochlorophyte grouping was suggested as useful in grouping those cyanobacteria with and without light-harvesting phycobilisomes, however it has become obvious that there have been multiple evolutionary origins of so-called prochlorophytes within widely different cyanobacterial taxonomic groups (Urbach *et al.*, 1992). Ribosomal RNA gene sequence analysis has become a more valid basis for establishing clearer evolutionary relationships (Urbach *et al.*, 1998; Honda *et al.*, 1999). Figure 2A shows one such analysis for 16S ribosomal RNA genes from a range of photosynthetic organisms. Cyanobacteria are clearly grouped within one radiation, with divisions within this primary radiation also being apparent. These analyses show them to be quite separate from β -proteobacteria, non-green algae, green algae, and higher plants. However, recent analysis of genomic sequences that have appeared for a number of cyanobacteria suggest that a division of cyanobacterial species based on their type of Rubisco may be much more appropriate (Badger *et al.*, 2002) particularly with respect to the evolution of their CCMs.

A phylogenetic tree for Rubisco from various photosynthetic bacteria is also shown in Fig. 2B. The photosynthetic organisms are grouped according to their Form 1 (L_8S_8) Rubisco large subunit types, originally described by Delwiche and colleagues (Delwiche, 1999), with Form 1A, B, C, and D groups shown. Cyanobacterial species are contained within both the Form 1A and 1B domains, as initially noted by Tabita and colleagues (Tabita, 1999). This variation of cyanobacteria is associated with the divergence between β -proteobacterial-like cyanobacteria such as *Prochlorococcus marinus* species and *Synechococcus* WH8102 and cyanobacteria such as *Synechocystis* PCC6803. This has led to a recent suggestion that two primary groupings of cyanobacteria can be established based on their Rubisco phylogeny (Badger *et al.*, 2002). These two groupings are referred to as α -cyanobacteria for those containing Form 1A Rubisco and β -cyanobacteria with Form 1B Rubisco.

The presence or absence of carboxysomes in various photosynthetic and chemoautotrophic bacterial species is also indicated in Fig. 2B. Carboxysomes are protein bodies which are surrounded by a protein shell and contain the Rubisco of the cell. Hence there is potential significance between the nature of the carboxysome structure and the type of Rubisco that it contains. Sequencing of the

α -cyanobacterial genomes has revealed that α -cyanobacteria possess carboxysomes that are significantly different from the carboxysomes found in β -cyanobacteria. Based on these observations it is suggested that these carboxysomes should be termed either α -carboxysomes for those found in Form 1A Rubisco-containing photosynthetic bacteria, including α -cyanobacteria and proteobacteria such as *Thiobacillus* species (Shively *et al.*, 1998a, b) while β -carboxysomes are those associated with Form 1B Rubisco in the β -cyanobacteria (Price *et al.*, 1998). Little is known about the different physiological properties of α and β -carboxysomes or the potential ecological advantages that the possession of carboxysomes might confer. However, all cyanobacteria characterized to date have carboxysomes.

Carboxysome structure and phylogeny

Ideas about the structure and function of carboxysomes have been based on studies with experimental systems for both α and β -carboxysomes, although the α -carboxysomes studied have been those associated with proteobacteria such as *Thiobacillus* species rather than α -cyanobacteria (Cannon *et al.*, 2002). However, the majority of studies on carboxysome function have been centred on β -carboxysomes in model laboratory species of β -cyanobacteria that are commonly used.

The carboxysome is a protein microbody, consisting of a protein coat, analogous to that surrounding a virus, and an interior soluble protein phase containing most, if not all, of the cellular Rubisco in cyanobacteria (Beudeker *et al.*, 1980; Price *et al.*, 1992; McKay *et al.*, 1993). The Rubisco within these bodies appears to be packed into paracrystalline arrays (Shively *et al.*, 1973; Holthuijzen *et al.*, 1986). The composition of the protein shell has been most extensively characterized in α -carboxysomes from chemolithotrophic proteobacteria such as *Thiobacillus* (Cannon *et al.*, 2001). Studies with these carboxysomes have shown the shell to consist of at least four different types of polypeptides. A number of small polypeptides (8–12 kDa), all related to each other, have been identified and named CsoS1, peptide A and peptide B. Frequently there are several members of each polypeptide type. Homologues of these peptides have been identified in β -cyanobacterial genomes and include CcmK, L, and O. Together, these proteins are related to each other by coding for proteins containing one or more regions of homology to bacterial micro-compartment domains (pfam 00936). These conserved structural domains have been identified by comparison of CcmK, CcmL, CcmO, or CsoS1-like proteins involved in carboxysome formation in α and β -cyanobacteria and β -proteobacteria (Price *et al.*, 1998; Shively *et al.*, 1998a, b; Cannon *et al.*, 2001) as well as more recently discovered genes associated with enteric proteobacteria containing carboxysome-like micro-com-

partments specialized in both propanediol and ethanolamine metabolism and detoxification (Bobick *et al.*, 1999; Kofoed *et al.*, 1999). Figure 3 shows a phylogenetic tree highlighting the relationships of the different polypeptides. CcmK, CcmO and CsoS1 proteins form separate groups while CcmL, peptideA and peptideB form another. Figure 3 clearly indicates the differences in the polypeptide composition between α and β -cyanobacteria. CcmK, L and O genes are contained in β -cyanobacterial genomes while CsoS1 and peptide A and B are found in α -cyanobacteria.

Carboxysome shells also appear to contain two other larger polypeptides that bear no homology to each other. In α -carboxysomes there are the CsoS2 (80–90 kDa) and CsoS3 (55–65 kDa), while in β -carboxysomes these appear to be replaced by CcmM (55–70 kDa) and CcmN (26 kDa) (Price *et al.*, 1998; Cannon *et al.*, 2001). In general, these proteins have no functional homologies in other bacterial systems, except that the CcmM protein has domains within it that are homologous to both γ -CA and to the small subunit of the Form 1B Rubisco protein (Price *et al.*, 1998; Ludwig *et al.*, 2000). The functional significance of these homologies is unknown.


For CO₂ fixation to occur in β -carboxysomes, it is envisaged that HCO₃⁻ diffuses through the proteinaceous shell of the carboxysome where a low activity of carbonic anhydrase inside the structure acts to catalyse the formation of CO₂ from HCO₃⁻ at rates high enough to saturate the carboxylation reaction of Rubisco. The role of the carboxysome as a site for CO₂ elevation was proposed by Reinhold *et al.* (1987, 1991). The models suggest that HCO₃⁻ diffuses into the carboxysome interior, and that CO₂ is generated by a specialized CA. CO₂ can be elevated within the carboxysome with the aid of some poorly understood diffusion barrier, such as the protein shell, that restricts CO₂ diffusion out of the carboxysome. Further information on the role of carboxysomes in the cyanobacterial CCM can be found in recent reviews (Price *et al.*, 1998; Kaplan and Reinhold, 1999; Cannon *et al.*, 2001).

Recent comparative genome analyses among different cyanobacteria (Cannon *et al.*, 2001, 2002; Hess *et al.*, 2001; Badger *et al.*, 2002) and the analysis in Fig. 3 have highlighted for the first time that α -cyanobacteria have α -carboxysomes rather than the β -carboxysomes previously studied in β -cyanobacteria. This suggests that these two carboxysome types have either been inherited or evolved in parallel, in association with the Form 1A and Form 1B Rubiscos found in each cyanobacterial group. Badger *et al.* (2002) have suggested the possibility that both Rubisco and carboxysome genes could be inherited by a single lateral gene transfer event as in many cyanobacteria they are often found organized on contiguous regions of the chromosome (Badger *et al.*, 2002).

The presence of a specific carboxysomal CA enzyme is interesting in that such a CA gene product has only been

identified in β -carboxysomes from a number of β -cyanobacteria (Fukuzawa *et al.*, 1992; Price *et al.*, 1992; Yu *et al.*, 1992). The emergence of complete genome sequences for a number of α and β -cyanobacteria has begun to confuse this picture. None of the α -cyanobacteria sequenced so far has a recognizable carboxysomal β -CA homologue, although one beta-CA (note that nomenclature is not related to α and β terminology used for carboxy-

somes and cyanobacteria) is present in the *Synechococcus* WH8102 genome (Badger *et al.*, 2002). In addition, the recently published genomes for the β -cyanobacteria *Trichodesmium erythraeum* (<http://genome.ornl.gov/microbial/tery/>) and *Thermosynechococcus elongatus* (<http://www.kazusa.or.jp/cyanobase/Thermo/index.html>) have also indicated a lack of any recognizable CA genes. The absence of any clearly identifiable CA genes in either α or



β-cyanobacteria is intriguing and points to the potential for a different mode of carboxysome function in these cyanobacteria.

Carbonic anhydrases

An analysis of possible alpha, beta and gamma carbonic anhydrases (Smith and Ferry, 2000) in the cyanobacterial genomes shows that there is a wide diversity in carbonic anhydrase gene content (Badger *et al.*, 2002). As noted above, a β-carboxysomal CA gene is present in many but not all β-cyanobacteria. In addition to this, one or more other beta-CAs may also be present, including *ecaB* (So and Espie, 1998). The β-cyanobacteria may also possess an alpha-CA (Soltes-Rak *et al.*, 1997). There are no clearly proven gamma-CAs in any of the cyanobacteria, although the CcmM protein in β-cyanobacteria has an amino terminal domain that could potentially contain an alpha-CA active site (Ludwig *et al.*, 2000) and ferripyochelin has some homology to gamma CA enzymes (Smith and Ferry, 2000). The absence of any identifiable CA genes in the α-cyanobacteria and at least two β-cyanobacterial genomes remains to be explained.

Ci transporters

Regardless of which form of Ci is presented to the cell, CO₂ or HCO₃⁻, the available evidence indicates that HCO₃⁻ is the species accumulated within the bulk cytoplasm. Furthermore, HCO₃⁻ and CO₂ are only slowly interconverted through the apparent absence of carbonic anhydrase activity in the general cytosol (Volokita *et al.*, 1984; Price and Badger, 1989). Being an ionic form of Ci, HCO₃⁻ is

much less permeable to lipid membranes than the uncharged CO₂ molecule and only slowly leaks from the cell, unless leakage is facilitated through a channel. The best support for the view that HCO₃⁻ is the accumulated species is that ectopic expression of human carbonic anhydrase within the cytoplasm of *Synechococcus* PCC7942 cells leads to a debilitating leakage of CO₂ from the cells due to rapid equilibration between HCO₃⁻ and CO₂ in the cytosol (Price and Badger, 1989).

Current evidence indicates that there are 4–5 modes of Ci uptake identified from research with *Synechococcus* PCC7942 and *Synechocystis* PCC6803 (Price *et al.*, 2002; Shibata *et al.*, 2002a). The four modes so far identified are discussed in more detail in Figs 4 and 5, but briefly they are: (1) BCT1, an inducible high affinity HCO₃⁻ transporter encoded by *cmpABCD* and belonging to the bacterial ATP binding cassette (ABC) transporter family (Omata *et al.*, 1999); (2) an inducible medium affinity Na⁺-dependent HCO₃⁻ transport system (Shibata *et al.*, 2002b); (3) a constitutive system for CO₂ uptake associated with a specialized form of a thylakoid-located NDH-1 complex, referred to as NDH-I₄ (Shibata *et al.*, 2001; Maeda *et al.*, 2002); and (4) a second specialized NDH-1 complex, named NDH-I₃ (Shibata *et al.*, 2001; Maeda *et al.*, 2002) that is inducible at limiting Ci conditions and exhibits a higher uptake affinity for CO₂.

The BCT1 HCO₃⁻ transporter

The high-affinity HCO₃⁻ transporter, BCT1, was the first cyanobacterial Ci transporter to have been conclusively identified and characterized. In *Synechococcus* PCC7942, BCT1 is encoded by the *cmpABCD* operon and is expressed under severe Ci limitation (Omata *et al.*,

Fig. 2. Phylogenetic trees for (A). 16S ribosomal small subunit DNA sequences; (B) Rubisco Form 1 (L₈S₈) large subunit protein sequences. Sequences are shown for photosynthetic and chemo-autotrophic bacteria, algae and embryophytes (including higher plants). Phylogenies are based on either DNA (A) or protein (B) sequences that were aligned using the program ClustalW version 1.7 (Thompson *et al.*, 1994). The phylogenetic trees are unrooted and were generated using the program PHYLIP version 3.6 alpha (J Felsenstein, University of Washington, Seattle, USA). All trees were generated using the Neighbor-Joining method, but they did not differ significantly from maximum likelihood parsimony analyses with respect to the conclusions drawn in the text. Ribosomal sequences were aligned using the online resources of the Ribosomal Database Project (<http://rdp.cme.msu.edu/html/>) (Maidak *et al.*, 2001) and analyses were based on 942 positions common to all sequences. Genbank accession numbers are as follows: *Trichodesmium erythraeum* (AF013030), *Nostoc punctiforme* (AF027655), *Anabaena variabilis* (AB016520), *Synechocystis* PCC6803 (D64000), *Synechococcus* PCC7942 (D88288), *Synechococcus* PCC6301 (X03538), *Synechococcus* WH8101 (AF001480), *Synechococcus* PCC7803 (AF081834), *Prochlorococcus* PAC1 (AF001471), *Prochlorococcus* MIT9313 (AF053399), *Synechococcus* PCC6307 (AF001477), *Chondrus crispus* (Z29521), *Cyanidium caldarium* (X52985), *Skeletonema costatum* (X82154), *Olisthodiscus luteus* (M82860), *Emiliana huxleyi* (X82156), *Ischrysis* sp. (X75518), *Chlamydomonas reinhardtii* (J01395), *Chlorella vulgaris* (D11347), *Coleochaete orbicularis* (U24579), *Marchantia polymorpha* (X04465), *Oryza sativa* (X15901), *Nicotiana tabacum* (U12813), *Pisum sativum* (X51598), *Rhodobacter sphaeroides* (D16425), *Rhodobacter capsulatus* (M34129), *Rhodopseudomonas palustris* (M59068), *Rhodospirillum rubrum* (M32020), *Ralstonia eutropha* (M32021), *Nitrosomonas europaea* (AF037106), *Thiobacillus thiooxidans* (M79401), *Thiobacillus ferro-oxidans* (X75266). Rubisco large subunit protein sequences references are as follows: *Synechococcus* WH8102, *Prochlorococcus* MIT9313, *Prochlorococcus* MED4, *Nostoc punctiforme*, *Trichodesmium erythraeum*, *Nitrosomonas europaea*, *Ralstonia metallidurans*, *Rhodobacter sphaeroides* I, *Rhodopseudomonas palustris* were retrieved from the DOE Joint Genome Project accessed through the Genome Channel (<http://compbio.ornl.gov/channel/>) and *Nostoc* PCC7120 from Cyanobase (<http://www.kazusa.or.jp/cyanobase/>). The PID numbers for other sequences were: *Synechococcus* PCC7803 (g1850939), *Pseudomonas hydrogothermophila* (g3183144), *Nitrobacter vulgaris* (g349304), *Thiobacillus ferro-oxidans* (g4836660), *Thiobacillus denitrificans* (g3183147), *Chromatium vinosum* (g131893), *Coleochaete nitellarium* (g5738990), *Phaeoceros laevis* (g12583955), *Triticum aestivum* (g132065), *Nicotiana tabacum* (g11841), *Marchantia polymorpha* (g131990), *Chlamydomonas reinhardtii* (g68154), *Cyanophora paradoxa* (g1352755), *Synechocystis* PCC6803 (g16331392), *Synechococcus* PCC6301 (g68159), *Ralstonia eutropha* (g6093905), *Rhodobacter sphaeroides* (g68170), *Xanthobacter flavus* (g68169), *Porphyridium aeruginum* (g730477), *Heterosigma akashiwo* (g11928), *Odontella sinensis* (g7436569), *Cyanidium caldarium* (g131941).




Fig. 3. A phylogenetic tree for CcmK-like proteins from α and β carboxysome-containing cyanobacteria and proteobacteria. Phylogenetic trees were constructed from aligning complete protein sequences as described in Fig. 2. Protein sequences for *Prochlorococcus* MED4 (peptide A, peptide B, CsoS1), *Prochlorococcus* MIT9313 (peptide A, peptide B, CsoS1), *Synechococcus* WH8102 (peptide B, 2 CsoS1 genes), *Nostoc punctiforme* (CcmO, 2 CcmK genes, CcmL) were retrieved from the DOE Joint Genome Project accessed through the Genome Channel (<http://compbio.ornl.gov/channel/>). Protein sequences for *Synechocystis* PCC6803 (CcmO, 4 CcmK genes, CcmL) and *Anabaena* PCC7120 (CcmO, 4 CcmK genes, CcmL) were retrieved from Cyanobase (<http://www.kazusa.or.jp/cyano/>). The PID numbers for other sequences were: *Synechococcus* PCC7002 CcmK (g3182943), CcmO (g2331051); *Synechococcus* PCC7942 ccmK (g416773), CcmO (g1176828), CcmL (g416774); *Thiobacillus neapolitanus* CsoS1 (g1169111, g1169109, g6014734), peptide A (g1176828).




Fig. 4. Models of two HCO_3^- transport systems identified in β -cyanobacteria. BCT1, the high-affinity traffic ATPase is shown on the left. SbtA (SLR1512), a potential $\text{Na}^+/\text{HCO}_3^-$ symport system is shown on the right. Further explanation about the operation of each system is described in the text.

1999). When the *cmp* operon was expressed under the control of a nitrate responsive promoter (*nirA*) the induced expression of the operon caused the appearance of high-affinity HCO_3^- uptake under high- CO_2 growth conditions (Omata *et al.*, 1999). BCT1, and its cyanobacterial orthologues, appear to be the first example of a primary transporter (uniporter) for HCO_3^- , but the transporter is clearly a member of the diverse subfamily of bacterial ABC transporters (Higgins, 2001). The *cmpA* gene codes for the precursor of the 42 kDa protein (CmpA) which has long been known to be induced under Ci limitation (Omata and Ogawa, 1986) and also under high-light stress (Reddy *et al.*, 1989). The protein sequence of CmpA is closely related to NrtA (Omata, 1991) which in turn has been confirmed as the nitrate/nitrite binding lipoprotein for the nitrate/nitrite transporter (*nrtABCD*) in *Synechococcus* sp. PCC7942 (Maeda and Omata, 1997). Recently, it has been shown that CmpA is indeed a binding protein that is specific for HCO_3^- with an affinity or K_D of around $5 \mu\text{M}$ (Maeda *et al.*, 2000).

Figure 4 shows a model for the operation of the BCT1 transporter. The *cmpABCD* genes code for four proteins:

the HCO₃⁻ binding protein (CmpA; 42 kDa); the intrinsic membrane protein (CmpB); a large extrinsic membrane protein with an consensus ATP binding site (CmpC), and a smaller related protein, CmpD, also with an ATP site. CmpB most probably forms a dimer within the membrane by analogy to other ABC transporters. The stoichiometry of CmpA to the other components of the transporter is apparently much greater than the intrinsic proteins of the transporter (Omata and Ogawa, 1986), but this is often a feature of binding proteins (Higgins, 2001). CmpA is a lipoprotein and it appears that CmpA proteins act as an array of substrate collectors for the transporter and are able to diffuse in two dimensions with the N-terminus of the mature polypeptide attached to the plasma membrane.

Comparative genomic analysis shows that the *cmpABCD* operon is found in freshwater β -cyanobacteria (see Fig. 6), but is absent from the marine *Synechococcus* PCC7002 (D Bryant, personal communication) and *Trichodesmium erythraeum* (see <http://genome.ornl.gov/microbial/tery/>) and is also absent from all three marine α -cyanobacteria species sequenced to date (Badger *et al.*, 2002).

Sodium-dependent HCO₃⁻ uptake

Cells of both *Synechococcus* PCC7942 and *Synechocystis* PCC6803 possess a Na⁺-dependent HCO₃⁻ uptake activity quite separate from BCT1 transport (Price *et al.*, 2002; Shibata *et al.*, 2002b). It has been suggested from physiological studies that cyanobacteria may possess a Na⁺/HCO₃⁻ symporter that is energized by a standing, inwardly directed Na⁺ gradient energized by a Na⁺/H⁺ antiporter (Espie and Kandasamy, 1994). Recently, Shibata *et al.* (2002a, b) have isolated a gene from *Synechocystis* PCC6803 that appears to code for Na⁺-dependent transport activity induced under Ci-limitation. The gene has been termed *sbtA* (SLR1512), and a mutant with lesions in *sbtA/cmpB/ndhD3/ndhD4* genes was unable to grow under low CO₂ at pH 7 or 9. This transporter is also depicted in Fig. 4. Analyses suggest that the predicted gene product of *sbtA* codes for a protein of 374 amino acids, and has potential for 8–10 membrane spanning domains. It also has a 60 amino acid hydrophilic domain near the centre of the protein that may represent a membrane extrinsic region, possibly facing the cytoplasm.




Fig. 5. A speculative model for the functioning of a specialized NDH-1_{3/4}-type complex to hydrate CO₂ to HCO₃⁻. The hydration is coupled to electron flow such as that supported by PSI cyclic electron transport. The model is based on the proposal of Price *et al.* (2002). It is postulated that ChpX and Y subunits are CO₂ hydration proteins bound to the cytoplasmic face of NDH-1_{3/4} complexes. The NdhF4/F3 and NdhD4/D3 subunits form part of the proton translocation channel. This specialized NDH-1 complex is proposed to drive the net hydration of CO₂ to HCO₃⁻ when coupled to photosynthetic electron transport. Here both NADPH and Fd_{red} are depicted as being potential electron donors to the NDH-1_{3/4} complexes. The energetics of CO₂ conversion may be further improved through the operation of a Q cycle similar to that operating in the *bf* complex. Equations 1–3 describe the reaction sequence proposed to occur in conventional CA enzymes (E) (Silverman, 2000; Smith *et al.*, 2000); the steps are (1) generation of a reactive hydroxyl group combined with spontaneous conversion of CO₂ to HCO₃⁻, (2) binding a water molecule to Zn and (3) abstraction of a proton to a nearby His residue and then conduction along a proton wire to a buffer molecule (B) in the bulk medium. The essential part of this proposal is that the last step (3a) be coupled to electron-driven translocation of protons to the thylakoid lumen, making hydroxyl-mediated net hydration of CO₂ largely irreversible in the light.




Fig. 6. Diversity in components of the cyanobacterial CCM. Depicted are four types of cyanobacterial cells, currently identifiable from genome sequencing projects. The groups are differentiated by the presence of either α or β -carboxysomes, and the complement of Ci transporters present. The two β -cyanobacterial groups are freshwater and marine types. For the marine α -cyanobacteria, there are two groups typified by either *Synechococcus* WH8102 or *Prochlorococcus* species. The species used to derive the groups are listed on the figure.

Strong homologues of the *sbtA* gene are present in many β -cyanobacterial genomes, however, only weak homologues appear to be present in the marine α -cyanobacteria (Badger *et al.*, 2002). Surprisingly, there are no homologues in the β -cyanobacterium *Thermosynechococcus elongatus* (<http://www.kazusa.or.jp/cyanobase/Thermo/>) and *Trichodesmium erythraeum* (<http://genome.ornl.gov/microbial/tery/>).

NDH-1 genes and CO₂ uptake

Previous studies have shown that the NDH-1 dehydrogenase complex is involved in enabling CO₂ uptake by cyanobacteria (Ogawa, 1992; Price *et al.*, 1998; Klughammer *et al.*, 1999; Ohkawa *et al.*, 2000a, b). However, within β -cyanobacterial species there may be a number of distinct types of NDH-1 complexes with different roles within the cell. The complete genome database for *Synechocystis* PCC6803 revealed that there are a number of NDH-1 genes in *Synechocystis* that are present as single copies in the genome. These are: *ndhAIGE*, *ndhB*, *ndhCJK*, *ndhH*, and *ndhL*. However, there are multiple copies of *ndhD* (6 homologues) and *ndhF* (3 homologues). Analysis of homology relationships indicated a large diversity in NdhD and NdhF proteins, with up to three groupings of NdhD being identifiable, as well as two groups of NdhF (Price *et al.*, 1998).

The NdhD1/D2 polypeptides together with NdhF1 probably are involved in forming a conventional respiratory NDH-1 complex, oxidizing NADPH and NADH and reducing plastoquinone, thus enabling cyclic electron transport around PSI (Ohkawa *et al.*, 2000a, b). The NdhD3/D4, together with NdhF3 and F4 components are suggested as components of two forms of a specialized NDH-1 complex involved in catalysing active CO₂ uptake by converting CO₂ to HCO₃⁻ within the cell (Ohkawa *et al.*, 2000a, b; Shibata *et al.*, 2001; Maeda *et al.*, 2002). The exact role of NdhD5/D6 polypeptides is unclear. In addition to this, two other genes/proteins are involved in enabling the CO₂ uptake activity of the NDH-1 complex, and these are referred to here as *chpX* and *chpY* (note that Shibata, Ogawa and colleagues have named these genes as *cupA* and *cupB*, while Price and co-workers have used *chpX* and *chpY* nomenclature; Shibata *et al.*, 2001, 2002a; Maeda *et al.*, 2002; Price *et al.*, 2002). Recent evidence from both Ogawa and Price's laboratories (Shibata *et al.*, 2001; Maeda *et al.*, 2002) have clearly indicated that the *ndhF3/ndhD3/chpY* genes code for polypeptides that are part of a high affinity CO₂ uptake NDH-1₃ complex, while the *ndhF4/ndhD4/chpX* genes code for a NDH-1₄ complex involved in low affinity CO₂ uptake.

Price *et al.* (2002) have speculated that the ChpX and ChpY polypeptides may be an integral part of the NDH-1 CO₂ uptake complex (NDH-1_{3/4}) and involved directly in the conversion of CO₂ to HCO₃⁻, linked to electron transport and proton translocation associated with the

NDH-1 complex. Recent attempts to determine the localization of these complexes suggest that they may be restricted to the thylakoid membrane (Ohkawa *et al.*, 2001) thus linking them directly to the photosynthetic electron transport chain. A model of the operation of such an NDH-1_{3/4} CO₂-uptake complex is shown in Fig. 5. Although it is apparent that the Chp proteins have no homologues with known CA protein families, multiple alignment of Chp proteins across ten α and β -cyanobacteria (Maeda *et al.*, 2002; Price *et al.*, 2002) show that it is possible to identify two conserved histidine residues and one conserved cysteine residue which could act as a potential Zn coordination site, as is found in conventional CA enzymes. The model shown in Fig. 5 is based on the theoretical potential of Chp proteins to bind Zn in a manner analogous to CA active sites, where Zn-OH⁻ can form and act as a strong nucleophile for CO₂ attack.

A potential reaction sequence for the conversion of CO₂ to HCO₃⁻ is shown in Fig. 5 and explained in the legend. Energization of CO₂ to HCO₃⁻ conversion would be achieved in a two-step process. Electron donation to the complex by donors such as NAD(P)H or Fd_{red} produces a reduced intermediate within the NDH-1 complex. This intermediate acts as a base (B) to abstract protons from the Zn-H₂O to produce the nucleophilic Zn-OH⁻ species. In the second step, the abstracted proton is translocated across the membrane to the lumen via a proton shuttle path within the hydrophobic proton channel subunits. By analogy with the *E. coli* NDH-1 complex (Friedrich and Scheide, 2000), the Ndh D and F subunits are part of the proton channel and would be specialized for their interactions with either the ChpX or ChpY subunits of each complex. The net result of these reactions is the hydration of CO₂ to HCO₃⁻ energized by electron transport through the NDH-1_{3/4} complexes. This model is still speculative and needs further evidence on the NDH-1 location of Chp proteins and their ability to bind Zn in the manner described above.

There is significant diversity in NDH-1_{3/4} gene content within and between α and β -cyanobacteria. The genomes of five β -cyanobacteria contain single copies of *ndhF*, *ndhD* and *chp* genes coding for both low and high-affinity CO₂ uptake (Badger *et al.*, 2002), however the marine β -cyanobacterium *Trichodesmium erythraeum* has only the low-affinity uptake forms. The marine α -cyanobacteria *Synechococcus* WH8102 has single copies of the *ndhF4*, *ndhD4* and *chpX* genes coding for a putative low-affinity CO₂ uptake system while the two *Prochlorococcus marinus* species have no homologues of either the low-affinity or high affinity NDH-1_{3/4} specific genes (Badger *et al.*, 2002). From what is known about β -cyanobacteria, this absence should mean that they lack the capacity for active CO₂ uptake, unless they possess another active CO₂ uptake system that is presently uncharacterized.

Diversity in CCM components between cyanobacteria and ecological adaptation.

Over the past two years, there has been a rapid increase in the number of complete cyanobacterial genome databases, and several more are due for completion in the next two years. This has provided the opportunity for comparative genomic analysis of the gene content and diversity of a range of evolutionarily distinct cyanobacteria from different environments. An analysis of existing genomes (Badger *et al.*, 2002), and the summaries outlined above, have shown that at present at least four types of cyanobacterial CCM strategies can be classified, defined on the presence of the types of carboxysomes present in the cells, and the complement of Ci transporters employed to accumulate HCO_3^- . A schematic summary of these four types is shown in the Fig. 6.

The presence of α and β -carboxysomes is perhaps the most striking variation between α and β -cyanobacteria. However, as nothing is known about the relative effectiveness of each type of carboxysome, it is difficult to speculate on the photosynthetic carboxylation advantages that each structure and its Rubisco may confer.

Most β -cyanobacterial genomes examined to date, except *Trichodesmium*, have genes correlated with both low and high affinity NDH-1 CO_2 uptake systems. This presumably implies that all these cyanobacteria are able to induce high affinity CO_2 uptake systems when inorganic carbon becomes limiting.

Common high and low-affinity HCO_3^- transport systems may also be present in most freshwater β -cyanobacteria, if *sbtA* (SLR1512) is a $\text{Na}^+/\text{HCO}_3^-$ transport gene, although *Thermosynechococcus elongatus* lacks an *sbtA* homologue. However, it should be noted that although *Synechocystis* contains the *cmpABCD* operon, and it is expressed (Omata *et al.*, 2001), there appears to be no associated high-affinity HCO_3^- uptake in this species (Shibata *et al.*, 2002b). The reason for this is unclear. The marine β -cyanobacterium *Synechococcus* PCC7002 lacks the *cmpABCD* operon (see NCBI unfinished bacterial genomes: <http://www.ncbi.nlm.nih.gov:80/cgi-bin/Entrez/framik?db=Genome&gi=5102>), however, it still manages to induce high-affinity HCO_3^- uptake when grown at low Ci (Sültemeyer *et al.*, 1995). There is a strong homologue of *sbtA* in this strain that may code for one of the HCO_3^- transport activities. Interestingly, the important N_2 -fixing marine β -cyanobacterium, *Trichodesmium erythraeum*, appears to lack both *cmpABCD* and *sbtA* and the molecular basis for any HCO_3^- uptake in this species remains to be determined.

The Ci transport genes of marine α -cyanobacteria may be quite different. *Synechococcus* WH8102 possesses genes that could code only for a CO_2 uptake system similar to the low-affinity CO_2 uptake system (NDH-1₄) from β -cyanobacteria. However, these genes are absent

from both *Prochlorococcus* species sequenced so far (Badger *et al.*, 2002). Observations with *Synechococcus* WH7803 (a close relative of WH8102) that the cells were able to evolve CO_2 during photosynthesis using HCO_3^- (Tchernov *et al.*, 1997) may be consistent with either the absence of CO_2 uptake genes in this species or the presence of only the low affinity uptake system. Extensive physiological studies of α -cyanobacterial species are lacking, but for *Prochlorococcus*, it would be expected that if these cyanobacteria do possess active HCO_3^- transport then they should evolve CO_2 during active photosynthesis. However, the nature of bicarbonate transport systems in α -cyanobacteria may be quite different from β -cyanobacteria. A strong homologue of *sbtA* appears to be absent (Badger *et al.*, 2002). If α and β -cyanobacteria diverged in their evolution prior to the development of bicarbonate transport systems, as carboxysome differences may suggest, then different types of HCO_3^- transport may have evolved independently.

The ecological significance of the differences in CCM gene content between α and β -cyanobacteria remains to be determined. However, it is clear that the marine α and β -cyanobacteria occupy quite different habitats compared with most freshwater β -cyanobacteria. The *Prochlorococcus* species, *Synechococcus* WH8102 and *Trichodesmium erythraeum* occur as dominant primary producers throughout oligotrophic oceanic waters (Moore *et al.*, 1998; Partensky *et al.*, 1999). In these environments it may be expected that inorganic carbon is never severely depleted and light and other nutrients may be major limiting factors. Many of the β -cyanobacteria occupy environments such as mats, films, estuarine situations, and alkaline lakes where higher population densities of organisms may occur, other nutrients may be more abundant and, overall, situations where inorganic carbon is a limiting resource may be much more common. This may explain why many β -cyanobacteria have the ability to induce various CO_2 and HCO_3^- transport systems as their environmental conditions change. However oceanic β -cyanobacteria such as *Trichodesmium* species that live in more oligotrophic waters are obviously different. The marine oceanic α -cyanobacteria may have developed a physiology where they may not have the ability to acquire or induce high affinity inorganic carbon transport systems, and in some species no active CO_2 uptake system may be present. Physiological studies of these species must be done to resolve this question.

Evolution of the cyanobacterial CCM

Considering current knowledge of cyanobacterial CCM genes and past evolutionary and climatic processes it is possible to speculate on the timing and evolution of CCMs in cyanobacteria. Figure 7 shows a chronological view of the development of photosynthetic cyanobacteria and




Fig. 7. The evolution of photosynthetic cyanobacteria, algae, higher plants, and chemoautotrophic proteobacteria over the past 3.5 billion years. (drawn with reference to Raven, 1997b, and Badger *et al.*, 2002). Also shown on the graph are the proposed changes in CO₂ and O₂ during the Phanerozoic period (Berner, 2001; Berner and Kothavala, 2001). CO₂ is shown as the ratio of past to present levels (R_{CO_2}) while O₂ is shown as percentage content. The shaded section indicates the period of CO₂ limitation combined with O₂ increase that may have initiated the development of CCMs in aquatic photosynthetic organisms.

algae over the past 3.5 billion years. This is plotted together with deduced changes in CO₂ and O₂ over the past 540 million years (the Phanerozoic era).

Past atmospheric CO₂ levels when cyanobacteria first arose were probably over 100-fold higher than present day conditions. This, combined with low O₂ conditions, would have meant that the original cyanobacteria would not have needed a CCM to achieve effective photosynthesis. The initial development of a CCM in cyanobacteria would have been triggered by changes in CO₂ and O₂ that caused CO₂ to be a limiting resource for photosynthesis and the Rubisco oxygenase reaction to become a significant problem. Clear records for changes in O₂ and CO₂ before about 600 million years ago are lacking, but it has been inferred that O₂ was near present levels by the beginning of the Phanerozoic and CO₂ may have been around 15–20 times current atmospheric levels. Given the properties of current cyanobacterial Rubiscos (Badger *et al.*, 1998), these enzymes should have been able to achieve efficient photosynthesis under these conditions. About 400 million years ago, there was a large decline in CO₂ levels and an almost doubling in the oxygen concentration. These changes would have placed significant pressures on both cyanobacterial and algal photosynthesis. It can be argued that this may have been the first time that major pressure was applied to photosynthetic organisms to develop CCMs (Raven, 1997a).

The first steps towards developing a cyanobacterial CCM may have been quite simple and speculation has been offered previously (Badger *et al.*, 2002) and is

outlined in Fig. 8. In the initial stages of CO₂ decline, the first step to developing a CCM would have been the evolution of a carboxysome structure for Rubisco. The cyanobacterial CCM is totally dependent on this structure and all other additions would have revolved around its presence. A carboxysome carbonic anhydrase would probably have been required at this stage as the rate of chemical conversion of HCO₃⁻ to CO₂ would have been too slow. As CO₂ limitation became more severe, the development of the NDH-1 based low and high-affinity CO₂ hydration process would have maintained adequate internal HCO₃⁻ pools and provided adequate CO₂ levels around Rubisco in the carboxysome. This process would have been based around the modification of an existing respiratory NDH-1 complex, and would have resulted in the efficient recycling of leaked CO₂ as well as net acquisition of CO₂ from outside the cell. Finally, as more extreme CO₂ limitation was imposed, the evolution of low and high affinity bicarbonate transport systems and high affinity CO₂-uptake would have been necessary.

Examining the genes involved in cyanobacterial carboxysomes and proteobacterial micro-compartments, it is obvious that the common components are the small *ccmK*, *cmmO*, *cmmL*-like and *csoS1*-bacterial micro-compartment genes (Fig. 3). The larger *csoS2* and *csoS3* genes are specific for Form 1A Rubisco α -carboxysomes, while the *cmmM* and *cmmN* genes are specific for Form 1B Rubisco β -carboxysomes. The view shown in Fig. 8 is that carboxysomes developed first in cyanobacteria and differentiated into both α and β carboxysomes. The appearance




Fig. 8. A speculative pathway for the evolution of the CCM and its components in α and β -cyanobacteria. (Drawn after Badger *et al.*, 2002.)

of carboxysomes in β -proteobacteria could have occurred through processes of lateral gene transfer which is becoming seen as a major part of bacterial evolution (Eisen, 2000; Brown *et al.*, 2001; Sicheritz-Ponten and Andersson, 2001).

The evolutionary divergence of α and β -cyanobacteria may have been a fairly ancient event and it has been argued that the types of cyanobacteria split before the advent of the primary endosymbiosis some 2 billion years ago (Tomitani *et al.*, 1999). If this is the case, then it could be argued that both cyanobacterial groups developed CCM mechanisms independently of each other rather than from a common ancestor having CCM components.

A polyphyletic origin of CCMs in cyanobacteria and algae

An independent and polyphyletic origin of CCM mechanisms in cyanobacteria, algae has been previously argued (Raven, 1997b; Badger *et al.*, 2002). A cornerstone to support this position has been the proposition that if CO_2 limitation were not imposed on cyanobacteria until the Phanerozoic, then this would strongly imply that the cyanobacteria that were the basis for the original primary endymbiotic event(s) probably did not have CCMs. Thus

the original Chlorophyte and Rhodophyte algae would also have lacked any CCM genetic elements in common with cyanobacteria, that could have aided their adaptation to falling CO_2 levels in the Phanerozoic. Chlorophytes and Rhodophytes as well as the secondary and tertiary endosymbiont algae that arose during the CO_2 -limitation of the Phanerozoic would all have needed to develop independent strategies for adapting to low CO_2 . Indeed, it has been suggested that the development of secondary endosymbiont algae may have been driven by this decline in CO_2 , as absorption into an acidic vacuolar structure may have made CO_2 more available by the conversion of HCO_3^- to CO_2 (Lee and Kugrens, 2000).

A search of the existing higher plant and algal DNA databases indicates there are no homologues of cyanobacterial CCM genes to be found. Thus carboxysome genes are almost exclusively restricted to cyanobacteria and some proteobacteria. Likewise, the NDH-1 $_{3/4}$ CO_2 uptake genes and the bicarbonate transport genes are restricted to cyanobacteria. An exception to this generalization is the occurrence of carboxysome structures in the plastids of Glaucocystophytes such as *Cyanophora paradoxa*. However, the evolutionary position of these algae has been very confusing. They retain a plastid that obviously resembles a cyanobacterium and if the plastids of red and green algae are assumed to be monophyletic then these algae appear to be an outgroup (Löffelhardt and Bohnert, 1994). Perhaps their plastids were obtained in a later endosymbiotic event involving a cyanobacterium that had developed a carboxysome structure. The algal genome databases are limited at present and are dominated by algal chloroplast genomes and *Chlamydomonas reinhardtii* ESTs, and it is possible that further sequencing of other algal species may uncover some CCM homologues in the nuclear genome. However, it would seem reasonable at this stage to assume that there were multiple origins of aquatic CCMs in algae and the ancestors of higher plants.

References

- Badger MR, Andrews TJ, Whitney SM, Ludwig M, Yellowlees DC, Leggat W, Price GD. 1998. The diversity and co-evolution of Rubisco, plastids, pyrenoids and chloroplast-based CCMs in the algae. *Canadian Journal of Botany* **76**, 1052–1071.
- Badger MR, Hanson DT, Price GD. 2002. Evolution and diversity of CO_2 concentrating mechanisms in cyanobacteria. *Functional Plant Biology* **29**, 407–416.
- Badger MR, Price GD. 1992. The CO_2 concentrating mechanism in cyanobacteria and microalgae. *Physiologia Plantarum* **84**, 606–615.
- Berner RA. 2001. Modeling atmospheric O_2 over Phanerozoic time. *Geochimica et Cosmochimica Acta* **65**, 685–694.
- Berner RA, Kothavala Z. 2001. GEOCARB III: A revised model of atmospheric CO_2 over phanerozoic time. *American Journal of Science* **301**, 182–204.
- Beudeker RF, Cannon GC, Kuenen JG, Shively JM. 1980. Relations between RuBP carboxylase, carboxysomes and CO_2

- fixing capacity in the obligate chemolithotroph *Thiobacillus neapolitana* grown under different limitations in the chemostat. *Archives of Microbiology* **1241**, 185–189.
- Bobick TA, Havemann GD, Busch RJ, Williams DS, Aldrich HC.** 1999. The propanediol utilization (*pdu*) operon of *Salmonella enterica* Serovar Typhimurium LT2 includes genes necessary for formation of polyhedral organelles involved in coenzyme B12-dependent 1,2-propanediol degradation. *Journal of Bacteriology* **181**, 5967–5975.
- Brown JR, Douady J, Italia MJ, Marshall WE, Stanhope J.** 2001. Universal trees based on large combined protein sequence data sets. *Nature Genetics* **28**, 281–285.
- Buick R.** 1992. The antiquity of oxygenic photosynthesis: evidence from stromatolites in sulphate-deficient archaean lakes. *Science* **255**, 74.
- Cannon GC, Bradburne CE, Aldrich HC, Baker SH, Heinhorst S, Shively JM.** 2001. Microcompartments in prokaryotes: carboxysomes and related polyhedra. *Applied and Environmental Microbiology* **67**, 5351–5361.
- Cannon GC, Heinhorst S, Bradburne CE, Shively JM.** 2002. Carboxysome genomics: a status report. *Functional Plant Biology* **29**, 175–182.
- Delwiche CF.** 1999. Tracing the thread of plastid diversity through the tapestry of life. *American Naturalist* **154**, S164–S177.
- Eisen JA.** 2000. Horizontal gene transfer among microbial genomes: new insights from complete genome analysis. *Current Opinion in Genetics and Development* **10**, 606–611.
- Espie GS, Kandasamy RA.** 1994. Monensin inhibition of Na⁺-dependent HCO₃⁻ transport distinguishes it from Na⁺-independent HCO₃⁻ transport and provides evidence for Na⁺/HCO₃⁻ symport in the cyanobacterium *Synechococcus* UTEX 625. *Plant Physiology* **104**, 1419–1428.
- Friedrich T, Scheide D.** 2000. The respiratory complex I of bacteria, archaea and eukarya and its module common with membrane-bound multisubunit hydrogenases. *FEBS Letters* **479**, 1–5.
- Fukuzawa H, Suzuki E, Komukai Y, Miyachi S.** 1992. A gene homologous to chloroplast carbonic anhydrase (*icfA*) is essential to photosynthetic carbon dioxide fixation by *Synechococcus* PCC7942. *Proceedings of the National Academy of Sciences, USA* **89**, 4437–4441.
- Hess W, Roca G, Ting C, Larimer F, Stilwagen S, Lamerdin J, Chisholm S.** 2001. The photosynthetic apparatus of *Prochlorococcus*: insights through comparative genomics. *Photosynthesis Research* (in press).
- Higgins CF.** 2001. ABC transporters: physiology, structure and mechanism—an overview. *Research in Microbiology* **152**, 205–210.
- Holthuijzen YA, Vanbreem JF, Konings WN, Vanbrugg EF.** 1986. Electron-microscopic studies of carboxysomes of *Thiobacillus neapolitanus*. *Archives of Microbiology* **144**, 258–262.
- Honda D, Yokota A, Sugiyama J.** 1999. Detection of seven major evolutionary lineages in cyanobacteria based on the 16S rRNA gene sequence analysis with new sequences of five marine *Synechococcus* strains. *Journal of Molecular Evolution* **48**, 723–739.
- Kaplan A, Reinhold L.** 1999. CO₂ concentrating mechanisms in photosynthetic microorganisms. *Annual Reviews in Plant Physiology and Plant Molecular Biology* **50**, 539–559.
- Klughammer B, Sültemeyer D, Badger MR, Price GD.** 1999. The involvement of NAD(P)H dehydrogenase subunits, NdhD3 and NdhF3, in high-affinity CO₂ uptake in *Synechococcus* sp. PCC7002 gives evidence for multiple NDH-I complexes with specific roles in cyanobacteria. *Molecular Microbiology* **32**, 1305–1315.
- Kofoid E, Rappleye C, Stojilkovic I, Roth J.** 1999. The 17-gene ethanolamine (eut) operon of *Salmonella typhimurium* encodes five homologues of carboxysome shell proteins. *Journal of Bacteriology* **181**, 5317–5329.
- Lee RE, Kugrens P.** 2000. Ancient atmospheric CO₂ and the timing of evolution of secondary endosymbioses. *Phycologia* **39**, 167–172.
- Löffelhardt W, Bohnert HJ.** 1994. Molecular biology of cyanobacteria. In: Bryant DA, ed. *The molecular biology of cyanobacteria*. Amsterdam: Kluwer, 65–89.
- Ludwig M, Sültemeyer D, Price GD.** 2000. Isolation of ccmKLMN genes from the marine cyanobacterium, *Synechococcus* sp. PCC7002 (Cyanobacteria), and evidence that CcmM is essential for carboxysome assembly. *Journal of Phycology* **36**, 1109–1118.
- Maeda S, Badger MR, Price GD.** 2002. Novel gene products associated with NdhD3/D4-containing NDH-I complexes are involved in photosynthetic CO₂ hydration in the cyanobacterium *Synechococcus* sp. PCC7942. *Molecular Microbiology* **43**, 425–435.
- Maeda S, Price GD, Badger MR, Enomoto C, Omata T.** 2000. Bicarbonate binding activity of the cmpA protein of the cyanobacterium *Synechococcus* PCC7942 is involved in active transport of bicarbonate. *Journal of Biological Chemistry* **275**, 20551–20555.
- Maeda SI, Omata T.** 1997. Substrate-binding lipoprotein of the cyanobacterium *Synechococcus* sp. strain PCC7942 involved in the transport of nitrate and nitrite. *Journal of Biological Chemistry* **272**, 3036–3041.
- Maidak BL, Cole JR, Lilburn TG, Parker CT, Saxman PR, Farris RJ, Garrity GM, Olsen GJ, Schmidt TM, Tiedje JM.** 2001. The RDP-II (Ribosomal Database Project). *Nucleic Acids Research* **29**, 173–174.
- McKay RML, Gibbs SP, Espie GS.** 1993. Effect of dissolved inorganic carbon on the expression of carboxysomes, localization of Rubisco and the mode of inorganic carbon transport in cells of the cyanobacterium *Synechococcus* UTEX 625. *Archives of Microbiology* **159**, 21–29.
- Moore LR, Roca G, Chisholm SW.** 1998. Physiology and molecular phylogeny of coexisting prochlorococcus ecotypes. *Nature* **393**, 464–467.
- Ogawa T.** 1992. Identification and characterization of the *icfA/ndhL* gene product essential to inorganic carbon transport of *Synechocystis* PCC6803. *Plant Physiology* **99**, 1604–1608.
- Ohkawa H, Pakrasi HB, Ogawa T.** 2000a. Two types of functionally distinct NAD(P)H dehydrogenases in *Synechocystis* sp strain PCC6803. *Journal of Biological Chemistry* **275**, 31630–31634.
- Ohkawa H, Price GD, Badger MR, Ogawa T.** 2000b. Mutation of *ndh* genes leads to inhibition of CO₂ uptake rather than HCO₃⁻ uptake in *Synechocystis* sp. strain PCC 6803. *Journal of Bacteriology* **182**, 2591–2596.
- Ohkawa H, Sonoda M, Shibata M, Ogawa T.** 2001. Localization of NAD(P)H dehydrogenase in the cyanobacterium *Synechocystis* sp. strain PCC6803. *Journal of Bacteriology* **183**, 4938–4939.
- Omata T.** 1991. Cloning and characterisation of the *nrtA* gene that encodes a 45 kDa protein involved in nitrate transport in the cyanobacterium *Synechococcus* PCC7942. *Plant and Cell Physiology* **32**, 151–157.
- Omata T, Gohta S, Takahashi Y, Harano Y, Maeda S.** 2001. Involvement of a CbbR homolog in low CO₂-induced activation of the bicarbonate transporter operon in cyanobacteria. *Journal of Bacteriology* **183**, 1891–1898.
- Omata T, Ogawa T.** 1986. Biosynthesis of a 42 kDa polypeptide in the cytoplasmic membrane of the cyanobacterium *Anacystis*

- nidulans* strain-R2 during adaptation to low CO₂ concentration. *Plant Physiology* **80**, 525–530.
- Omata T, Price GD, Badger MR, Okamura M, Gohta S, Ogawa T.** 1999. Identification of an ATP-binding cassette transporter involved in bicarbonate uptake in the cyanobacterium *Synechococcus* sp. strain PCC7942. *Proceedings of the National Academy of Sciences, USA* **96**, 13571–13576.
- Partensky F, Hess WR, Vaulot D.** 1999. *Prochlorococcus*, a marine photosynthetic prokaryote of global significance. *Microbiology and Molecular Biology Reviews* **63**, 106–127.
- Price GD, Badger MR.** 1989. Expression of human carbonic anhydrase in the cyanobacterium *Synechococcus* PCC7942 creates a high CO₂-requiring phenotype. Evidence for a central role for carboxysomes in the CO₂ concentrating mechanism. *Plant Physiology* **91**, 505–513.
- Price GD, Coleman JR, Badger MR.** 1992. Association of carbonic anhydrase activity with carboxysomes isolated from the cyanobacterium *Synechococcus* PCC7942. *Plant Physiology* **100**, 784–793.
- Price GD, Maeda S-I, Omata T, Badger MR.** 2002. Modes of inorganic carbon uptake in the cyanobacterium *Synechococcus* sp. PCC7942. *Functional Plant Biology* **29**, 131–149.
- Price GD, Sültemeyer D, Klughammer B, Ludwig M, Badger MR.** 1998. The functioning of the CO₂ concentrating mechanism in several cyanobacterial strains: a review of general physiological characteristics, genes, proteins and recent advances. *Canadian Journal of Botany* **76**, 973–1002.
- Raven JA.** 1997a. Putting the C back into phycology. *European Journal of Phycology* **32**, 319–333.
- Raven JA.** 1997b. The role of marine biota in the evolution of terrestrial biota—gases and genes—atmospheric composition and evolution of terrestrial biota. *Biogeochemistry* **39**, 139–164.
- Reddy KJ, Masamoto K, Sherman DM, Sherman LA.** 1989. DNA sequence and regulation of the gene (*cbpA*) encoding the 42 kilodalton cytoplasmic membrane carotenoprotein of the cyanobacterium *Synechococcus* sp. strain PCC7942. *Journal of Bacteriology* **171**, 3486–3493.
- Reinhold L, Kosloff R, Kaplan A.** 1991. A model for inorganic carbon fluxes and photosynthesis in cyanobacterial carboxysomes. *Canadian Journal of Botany* **69**, 984–988.
- Reinhold L, Zviman M, Kaplan A.** 1987. Inorganic carbon fluxes and photosynthesis in cyanobacteria—a quantitative model. In: Biggins J, ed. *Progress in photosynthesis research: proceedings of the VIIth international congress on photosynthesis*. Dordrecht: M Nijhoff Publishers, 289–296.
- Shibata M, Ohkawa H, Kaneko T, Fukuzawa H, Tabata S, Kaplan A, Ogawa T.** 2001. Distinct constitutive and low-CO₂-induced CO₂ uptake systems in cyanobacteria: genes involved and their phylogenetic relationship with homologous genes in other organisms. *Proceedings of the National Academy of Sciences, USA* **98**, 11789–11794.
- Shibata M, Ohkawa H, Katoh H, Shimoyama M, Ogawa T.** 2002a. Two CO₂-uptake systems: four systems for inorganic carbon acquisition in *Synechocystis* sp. strain PCC6803. *Functional Plant Biology* **29**, 123–129.
- Shibata M, Katoh H, Sonoda M, Ohkawa H, Shimoyama M, Fukuzawa H, Kaplan A, Ogawa T.** 2002b. Genes essential to sodium-dependent bicarbonate transport in cyanobacteria—function and phylogenetic analysis. *Journal of Biological Chemistry* **277**, 18658–18664.
- Shively JM, Ball FL, Kline BW.** 1973. Electron microscopy of the carboxysomes (polyhedral bodies) of *Thiobacillus neapolitanus*. *Journal of Bacteriology* **116**, 1405–1411.
- Shively JM, Bradburne CE, Aldrich HC, Bobick TA, Mehlmán JL, Jin S, Baker SH.** 1998a. Sequence homologues of the carboxysomal polypeptide CsoS1 of the thiobacilli are present in cyanobacteria and enteric bacteria that form carboxysomes—polyhedral bodies. *Canadian Journal of Botany* **76**, 906–916.
- Shively JM, Vankeulen G, Meijer WG.** 1998b. Something from almost nothing—carbon dioxide fixation in chemoautotrophs. *Annual Review of Microbiology* **52**, 191–230.
- Sicheritz-Ponten T, Andersson SGE.** 2001. A phylogenomic approach to microbial evolution. *Nucleic Acids Research* **29**, 545–552.
- Silverman DN.** 2000. Marcus rate theory applied to enzymatic proton transfer. *Biochimica et Biophysica Acta—Bioenergetics* **1458**, 88–103.
- Smith KS, Cosper NJ, Stalhandske C, Scott RA, Ferry JG.** 2000. Structural and kinetic characterization of an archaeal beta-class carbonic anhydrase. *Journal of Bacteriology* **182**, 6605–6613.
- Smith KS, Ferry JG.** 2000. Prokaryotic carbonic anhydrases. *FEMS Microbiology Reviews* **24**, 335–366.
- So AKC, Espie GS.** 1998. Cloning, characterization and expression of carbonic anhydrase from the cyanobacterium *Synechocystis* PCC6803. *Plant Molecular Biology* **37**, 205–215.
- Soltes-Rak E, Mulligan ME, Coleman JR.** 1997. Identification and characterization of a gene encoding a vertebrate-type carbonic anhydrase in cyanobacteria. *Journal of Bacteriology* **179**, 769–774.
- Sültemeyer D, Price GD, Yu JW, Badger MR.** 1995. Characterisation of carbon dioxide and bicarbonate transport during steady-state photosynthesis in the marine cyanobacterium *Synechococcus* strain PCC7002. *Planta* **197**, 597–607.
- Tabita FR.** 1999. Microbial ribulose 1,5-bisphosphate carboxylase/oxygenase: a different perspective. *Photosynthesis Research* **60**, 1–28.
- Tchernov D, Hassidim M, Luz B, Sukenik A, Reinhold L, Kaplan A.** 1997. Sustained net CO₂ evolution during photosynthesis by marine microorganisms. *Current Biology* **7**, 723–728.
- Thompson JD, Higgins HD, Gibson TL.** 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**, 4673–4680.
- Tomitani A, Okada K, Miyashita H, Matthijs HCP, Ohno T, Tanaka A.** 1999. Chlorophyll *b* and phycobilins in the common ancestor of cyanobacteria and chloroplasts. *Nature* **400**, 159–162.
- Urbach E, Robertson DL, Chisholm SW.** 1992. Multiple evolutionary origins of prochlorophytes within the cyanobacterial radiation. *Nature* **355**, 267–270.
- Urbach E, Scanlan DJ, Distel DL, Waterbury JB, Chisholm SW.** 1998. Rapid diversification of marine picophytoplankton with dissimilar light-harvesting structures inferred from sequences of *Prochlorococcus* and *Synechococcus* (cyanobacteria). *Journal of Molecular Evolution* **46**, 188–201.
- Volokita M, Zenvirth D, Kaplan A, Reinhold L.** 1984. Nature of the inorganic carbon species actively taken up by the cyanobacterium *Anabaena variabilis*. *Plant Physiology* **76**, 599–602.
- Yu JW, Price GD, Song L, Badger MR.** 1992. Isolation of a putative carboxysomal carbonic anhydrase gene from the cyanobacterium *Synechococcus* PCC7942. *Plant Physiology* **100**, 794–800.